

Further Pure 1 - January 2009

(1) $y_{n+1} = y_n + hf(x_n)$

$$x_1 = 0$$

$$y_1 = 1$$

$$h = 0.2$$

$$y_2 = 1 + 0.2 \times \sqrt{1 + 0^2} \\ = 1.2$$

$$x_2 = 0.2$$

$$y_2 = 1.2$$

$$h = 0.2$$

$$y_3 = 1.2 + 0.2 \times \sqrt{1 + 0.2^2} \\ = 1.40396 \quad (5dp)$$

(2) a) $2 - 3i$ (conjugate)

b) $x^2 + \boxed{\text{sum}}x + \boxed{\text{product}} = 0$

$$\boxed{\text{sum}} \quad 2 + 3i + 2 - 3i = 4 \quad \rightarrow b = -4$$

$$\boxed{\text{product}} \quad (2 + 3i)(2 - 3i) = 4 + 9 = 13 \quad \rightarrow c = 13$$

(3) $\tan: \theta = n\pi + \alpha$

$$\text{key angle } (\alpha) = \tan^{-1}(\sqrt{3}) = \pi/3$$

$$\rightarrow \pi/2 - 3x = n\pi + \pi/3$$

$$-3x = n\pi - \pi/6$$

$$\rightarrow x = -\pi/3 \cdot \pi + \pi/18$$

which is same as $x = \pi/3 \cdot \pi + \pi/18$

(4) a) $\sum r = \frac{1}{2} n(n+1)$

$$\sum r^2 = \frac{1}{6} n(n+1)(2n+1)$$

$$\rightarrow 3 \sum r^2 + 3 \sum r + \sum 1$$

$$= \frac{1}{2} n(n+1)(2n+1) + \frac{3}{2} n(n+1) + n$$

$$= \frac{1}{2} n \left[(n+1)(2n+1) + 3(n+1) + 2 \right]$$

$$= \frac{1}{2} n \left[2n^2 + 3n + 1 + 3n + 3 + 2 \right]$$

$$= \frac{1}{2} n \left[2n^2 \right] = n^3$$

$$b) \sum_{n=1}^{2n} = \sum_{n=1}^{2n} - \sum_{n=1}^n = (2n)^3 - (n)^3 = 7n^3$$

$$③ a) i) A+B = \begin{bmatrix} k & k \\ k & -k \end{bmatrix} + \begin{bmatrix} -k & k \\ k & k \end{bmatrix} = \begin{bmatrix} 0 & 2k \\ 2k & 0 \end{bmatrix}$$

$$ii) A^2 = \begin{bmatrix} k & k \\ k & -k \end{bmatrix} \begin{bmatrix} k & k \\ k & -k \end{bmatrix} = \begin{bmatrix} k & k \\ k & -k \end{bmatrix} \begin{bmatrix} 2k^2 & 0 \\ 0 & 2k^2 \end{bmatrix}$$

$$b) (A+B)^2 = \begin{bmatrix} 0 & 2k \\ 2k & 0 \end{bmatrix} \begin{bmatrix} 0 & 2k \\ 2k & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 2k \\ 2k & 0 \end{bmatrix} \begin{bmatrix} 4k^2 & 0 \\ 0 & 4k^2 \end{bmatrix}$$

$$B^2 = \begin{bmatrix} -k & k \\ k & k \end{bmatrix} \begin{bmatrix} -k & k \\ k & k \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 2k^2 & 0 \\ 0 & 2k^2 \end{bmatrix}$$

$$= \begin{bmatrix} -k & k \\ k & k \end{bmatrix} \begin{bmatrix} 2k^2 & 0 \\ 0 & 2k^2 \end{bmatrix}$$

$$\therefore A^2 + B^2 = \begin{bmatrix} 2k^2 & 0 \\ 0 & 2k^2 \end{bmatrix} + \begin{bmatrix} 2k^2 & 0 \\ 0 & 2k^2 \end{bmatrix} = \begin{bmatrix} 4k^2 & 0 \\ 0 & 4k^2 \end{bmatrix} = (A+B)^2$$

$$c) i) A^2 = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \text{Enlargement, scale factor 2, centre } (0,0)$$

$$ii) A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \text{ Enlargement SF must } = \sqrt{2}$$

$$\rightarrow \sqrt{2} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix}$$

Reflection

$$\cos(2\theta) = \frac{1}{\sqrt{2}}$$

$$\rightarrow 2\theta = 45^\circ$$

$$\rightarrow \theta = 22.5^\circ$$

\therefore Reflection in line $y = \tan(22.5^\circ)x$

(6) a) i) Denominator = 0 when $x = 0$ or $x = 2$

$$\rightarrow x = 0, x = 2$$


As $x \rightarrow \infty, y \rightarrow 1/2 \rightarrow y = 1/2$

$$\text{ii) } y = 0 \rightarrow 0 = \frac{(x-1)(x-3)}{x(x-2)}$$

$$\rightarrow 0 = (x-1)(x-3)$$


$$\rightarrow x = 1 \text{ or } x = 3 \quad \text{or } (1,0) \text{ or } (3,0)$$

iii)




b) curve below $x = \text{axis}$ when: $0 < x < 1$ and $2 < x < 3$

(7) a)



$c = d$



$$\frac{r-a}{c} = \frac{b-a}{c-d}$$

$$r-a = c \left(\frac{b-a}{c-d} \right)$$

$$\rightarrow r = a + c \left(\frac{b-a}{c-d} \right)$$

$$b) i) a = 2 \rightarrow c = 20(2) - (2)^4 = 24$$

$$b = 3 \rightarrow d = 20(3) - 3^4 = -21$$

$$\rightarrow r = 24 \left(\frac{3 - 3}{24 - (-21)} \right) + 2 = 38/15$$

$$ii) \text{ at } \beta, f(x) = 0$$

$$\rightarrow 20x - x^4 = 0$$

$$\rightarrow x^4 - 20x = 0$$

$$x(x^3 - 20) = 0$$

$$x = 0$$

This is not β

$$x^3 - 20 = 0$$

$$\rightarrow x = \sqrt[3]{20} = \beta$$

$$\therefore \beta - r = \sqrt[3]{20} - 38/15 = 0.181...$$

$$(8) a) \int_1^{\infty} x^{-3/4} dx = \int_1^n x^{-3/4} dx$$

$$= \left[4x^{1/4} \right]_1^n = 4\sqrt[4]{n} - 4\sqrt[4]{1}$$

As $n \rightarrow \infty$, $4\sqrt[4]{n}$ does not converge to a limit, so \int has no value

$$b) \int_1^{\infty} x^{-5/4} dx = \int_1^n x^{-5/4} dx = \left[-4x^{-1/4} \right]_1^n$$

$$= \frac{-4}{\sqrt[4]{n}} - \left(\frac{-4}{1} \right)$$


$$\text{As } n \rightarrow \infty, \frac{-4}{\sqrt[4]{n}} \rightarrow 0, \text{ and } \therefore \int \rightarrow 4$$

c) First component still has no value, so subtraction 4 makes no difference. = NO VALUE.

$$(9) a) \text{ From formula book, asymptotes } = \frac{y}{1} = \pm \frac{y}{\sqrt{2}}$$

$$\rightarrow y = \sqrt{2}x \text{ and } y = -\sqrt{2}x$$

b)



$$i) \text{ if } y = x + c \quad \rightarrow \quad x^2 - \frac{(x+c)^2}{2} = 1$$

$$\rightarrow 2x^2 - (x+c)^2 = 2$$

$$2x^2 - x^2 - 2xc - c^2 = 2$$

$$x^2 - 2cx - (c^2 + 2) = 0$$

$$ii) \text{ Discriminant: } b^2 - 4ac$$

$$= (-2c)^2 - 4 \times 1 \times -(c^2 + 2)$$

$$= 4c^2 + 4c^2 + 8$$

$$= 8c^2 + 8$$

Whatever the value of c , this is always +ve, so
2 distinct solutions

iii) Solve quadratic:

$$x = \frac{2c \pm \sqrt{8c^2 + 8}}{2}$$

$$x = \frac{2c \pm \sqrt{4} \sqrt{2c^2 + 2}}{2}$$

$$x = c \pm \sqrt{2c^2 + 2}$$

$$y = x + c$$

$$\rightarrow y = 2c \pm \sqrt{2c^2 + 2}$$