

# FP1 - January 2010

$$\textcircled{1} \text{ a) } \alpha + \beta = \frac{6}{3} = 2$$
$$\alpha\beta = \frac{1}{3}$$

$$\text{b) } \alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$$
$$= 2^3 - 3 \times \frac{1}{3} \times 2 = 6$$

$$\text{c) } \boxed{\text{Sum}} \quad \alpha^2/\beta + \beta^2/\alpha = \frac{\alpha^3}{\alpha\beta} + \frac{\beta^3}{\alpha\beta}$$
$$= \frac{\alpha^3 + \beta^3}{\alpha\beta} = \frac{6}{1/3} = 18$$

$$\boxed{\text{Product}} \quad \alpha^2/\beta \times \beta^2/\alpha = \frac{\alpha^2\beta^2}{\alpha\beta} = \alpha\beta = \frac{1}{3}$$

$$\rightarrow x^2 - \boxed{\text{Sum}}x + \boxed{\text{Product}} = 0$$

$$\rightarrow x^2 - 18x + \frac{1}{3} = 0$$

$$\rightarrow 3x^2 - 54x + 1 = 0$$

$$\textcircled{2} \text{ a) } z^2 = (1+i)(1+i) = 1 + 2i + i^2$$
$$= 1 + 2i - 1 = 2i$$

$$\text{b) } z^8 = (z^2)^4 = (2i)^4 = 2^4 \times i^4$$
$$= 16 \times (-1)(-1) = 16$$

$$\text{c) } (z^*)^2 = (1-i)(1-i) = 1 - 2i + i^2 = -2i$$
$$= -z^2$$

$$(3) \quad \theta = 2n\pi + a, \quad \theta = 2n\pi + (\pi - a)$$

Key angle:  $\sin^{-1}(1) = \pi/2$

$$4x + \pi/4 = 2n\pi + \pi/2, \quad 4x + \pi/4 = 2n\pi + \pi/2$$

SAME!

$$\rightarrow 4x = 2n\pi + \pi/4$$

$$\rightarrow x = \frac{1}{2}n\pi + \pi/16$$

$$(4) \quad a) \quad A - I = \begin{bmatrix} 1 & 4 \\ 3 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 4 \\ 3 & 0 \end{bmatrix}$$

$$(A - I)^2 = \begin{bmatrix} 0 & 4 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 0 & 4 \\ 3 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 4 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 12 & 0 \\ 0 & 12 \end{bmatrix} = 12I$$

$$b) \quad A - B = \begin{bmatrix} 1 & 4 \\ 3 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 3 \\ p & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 3-p & 0 \end{bmatrix}$$

$$(A - B)^2 = \begin{bmatrix} 0 & 1 \\ 3-p & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 3-p & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 3-p & 0 \end{bmatrix} \begin{bmatrix} 3-p & 0 \\ 0 & 3-p \end{bmatrix}$$

$$3-p = 12$$

$$\rightarrow p = -9$$

(5) a) The integral is undefined when  $x=0$

As  $x \rightarrow 0$ ,  $1/\sqrt{x} \rightarrow \infty$

$$\begin{aligned} \text{b) i) } \int_0^{1/16} x^{-1/2} &= \int_p^{1/16} x^{-1/2} \\ &= \left[ 2x^{1/2} \right]_p^{1/16} = \left[ 2(1/16)^{1/2} - 2p^{1/2} \right] \\ &= 1/2 - \cancel{2\sqrt{p}} \end{aligned}$$

As  $p \rightarrow 0$ ,  $2\sqrt{p} \rightarrow 0$ , so  $\int \rightarrow 1/2$

$$\begin{aligned} \text{ii) } \int_0^{1/16} x^{-5/4} &= \int_p^{1/16} x^{-5/4} \\ &= \left[ -4x^{-1/4} \right]_p^{1/16} = \left[ -4/\sqrt[4]{x} \right]_p^{1/16} \\ &= -4/\sqrt[4]{1/16} - -4/\sqrt[4]{p} \end{aligned}$$

As  $p \rightarrow 0$ ,  $4/\sqrt[4]{p} \rightarrow \infty$ , so integral has no finite value.

$$\text{(6) a) i) } \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 3 & 3 & 1 \\ 1 & 1 & 2 & 2 \\ 3 & a & a & 3 \\ 2 & 2 & 4 & 4 \end{bmatrix} \rightarrow \begin{matrix} (3,2) & (9,2) \\ (9,4) & (3,4) \end{matrix}$$

ii) See Mark Scheme

b) i) See Mark Scheme.

ii) Rotation  $90^\circ$  cw =  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$




c)  $R_1 \rightarrow R_3$  must multiply in reverse order

$$\rightarrow \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$$

$$\begin{array}{c|c} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} & \begin{bmatrix} 0 & 2 \\ -3 & 0 \end{bmatrix} \\ \hline & \end{array}$$

⑦ a) i)  $x=2$  As  $x \rightarrow \infty, y \rightarrow \frac{1}{\infty} = 0$   
 $\rightarrow y=0$



b)  $x-3 = \frac{1}{(x-2)^2}$

$$\rightarrow (x-3)(x-2)^2 = 1$$

$$\rightarrow (x-3)(x-2)^2 - 1 = 0$$

$$f(x) = (x-3)(x-2)^2 - 1$$

$$f(3) = -1$$

$$f(4) = 3$$

Sign change,  $\therefore$  root lies between 3 & 4

ii)	Interval	Midpoint $(x)$	Value of $f(x)$	
	3 - 4	3.5	0.125 <del>more 55555...</del>	→ { Between 3 & 3.5 }
	3 - 3.5	3.25	-0.984375	
	∴ must be between 3.25 & 3.5			

8) a)  $\sum r^3 + \sum r$   
 $= \frac{1}{4} n^2 (n+1)^2 + \frac{1}{2} n (n+1)$   
 $= \frac{1}{4} n (n+1) + [n(n+1) + 2]$   
 $= \frac{1}{4} n (n+1) + [n^2 + n + 2]$

b)  $8 \sum r^2 = \frac{8}{6} n (n+1) (2n+1)$

→  $\frac{1}{4} n (n+1) (n^2 + n + 2) = \frac{8}{6} n (n+1) (2n+1)$

$\div n(n+1)$  →  $\frac{1}{4} (n^2 + n + 2) = \frac{8}{6} (2n+1)$

$\times 12$  →  $3(n^2 + n + 2) = 16(2n+1)$

→  $3n^2 + 3n + 6 = 32n + 16$

→  $3n^2 - 29n - 10 = 0$

→  $(3n+1)(n-10) = 0$

↓                      ↓

$n = -1/3$

$n = 10$

$n$  must be a positive integer →  $n = 10$

9) a) For  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  asymptotes are  $\frac{x}{a} = \pm \frac{y}{b}$

when  $x=2, y=0$

→  $\frac{4}{a^2} - 0 = 1 \rightarrow a^2 = 4 \rightarrow a = 2$

Asymptote =  $y = 2x \rightarrow y = \frac{bx}{a}$  same as  $y = 2x$

→  $\frac{b}{a} = 2 \rightarrow \frac{b}{2} = 2$

→  $b = 4$

b) Equation of line:  $y - 0 = m(x - 1)$

$$\rightarrow y = mx - m$$

Intersects at:

$$\frac{x^2}{4} - \frac{(mx - m)^2}{16} = 1$$

$$\rightarrow 4x^2 - \cancel{m^2 x^2} (mx - m)^2 = 16$$

$$\rightarrow 4x^2 - m^2 x^2 + 2m^2 x - m^2 = 16$$

$$0 = m^2 x^2 - 4x^2 - 2m^2 x + m^2 + 16$$

$$\rightarrow (m^2 - 4)x^2 - (2m^2)x + (m^2 + 16) = 0$$

c) For equal roots,  $b^2 - 4ac = 0$

$$\rightarrow (-2m^2)^2 - 4 \times (m^2 - 4)(m^2 + 16) = 0$$

$$\rightarrow 4m^4 - 4[m^4 + 12m^2 - 64] = 0$$

$$\rightarrow m^4 - m^4 - 12m^2 + 64 = 0$$

$$\rightarrow -12m^2 + 64 = 0$$

$$\rightarrow 12m^2 = 64 \rightarrow 3m^2 = 16$$

d)  $3m^2 = 16 \rightarrow m^2 = 16/3$

Find x-co-ord in:  $(m^2 - 4)x^2 - (2m^2)x + (m^2 + 16) = 0$

$$\rightarrow (16/3 - 4)x^2 - 2(16/3)x + 16/3 + 16 = 0$$

$$\rightarrow 4/3 x^2 - 32/3 x + 16/3 + 16 = 0$$

$$\boxed{\times 3} \rightarrow 4x^2 - 32x + 64 = 0$$

$$\rightarrow x^2 - 8x + 16 = 0$$

$$(x - 4)(x - 4) = 0$$

$$\rightarrow x = 4$$

Need to find y:

$$x^2/2^2 - y^2/4^2 = 1$$

$$\rightarrow 4^2/2^2 - y^2/4^2 = 1$$

$$\rightarrow 4 - y^2/16 = 1$$

$$64 - y^2 = 16$$

$$y^2 = 48$$

$$\rightarrow y = \pm \sqrt{48}$$

$$\rightarrow y = \pm 4\sqrt{3}$$

$$(4, 4\sqrt{3}) \quad \text{AND} \quad (4, -4\sqrt{3})$$