

### Further Puc 1 - June 2013

$$(1) \quad x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$x_1 = 10$$

$$f(x) = x^3 - x^2 + 4x - 900$$

$$f(10) = 10^3 - 10^2 + 4(10) - 900 = 40$$

$$f'(x) = 3x^2 - 2x + 4$$

$$f'(10) = 3(10)^2 - 2(10) + 4 = 284$$

$$\rightarrow x_{n+1} = 10 - \frac{40}{284} = 9.85915... \\ = 9.859 \text{ (4 s.d.)}$$

$$(2) \text{ a) i) } \begin{bmatrix} p & 2 \\ 4 & p \end{bmatrix} - \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} p-3 & 1 \\ 2 & p-3 \end{bmatrix}$$

$$\text{ii) } \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix}$$

$$\begin{bmatrix} p & 2 \\ 4 & p \end{bmatrix} \begin{bmatrix} 3p+4 & p+6 \\ 12+2p & 4+3p \end{bmatrix} \rightarrow \begin{bmatrix} 3p+4 & p+6 \\ 12+2p & 4+3p \end{bmatrix}$$

$$\text{b) } A - B + AB = \begin{bmatrix} p-3 & 1 \\ 2 & p-3 \end{bmatrix} + \begin{bmatrix} 3p+4 & p+6 \\ 12+2p & 4+3p \end{bmatrix}$$

$$= \begin{bmatrix} 4p+1 & p+7 \\ 14+2p & 1+4p \end{bmatrix} \quad \text{needs to be } K \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\rightarrow p+7 = 0 \quad \rightarrow p = -7$$

$$\rightarrow \begin{bmatrix} -27 & 0 \\ 0 & -27 \end{bmatrix} = -27 I \quad K = -27$$

$$(3) a) \cos: \theta = 360n \pm \alpha$$

$$\text{Key angle} = \cos^{-1}(\cos 65) = 65 = \alpha$$

$$\rightarrow 5x + 40 = 360n \pm 65$$

$$\rightarrow 5x = 360n - 40 \pm 65$$

$$\rightarrow x = 72n - 8 \pm 13$$

$$\rightarrow x = 72n - 5, \quad x = 72n - 21$$

$$b) \cos(\pi/4) = 1/\sqrt{2}$$

$$\sin(\pi/12) = \frac{\sqrt{3} - 1}{2\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}} \left( \frac{\sqrt{3} - 1}{2} \right)$$

$$= \frac{1}{\sqrt{2}} \left( \frac{\sqrt{3}}{2} - \frac{1}{2} \right)$$

$$\begin{array}{ccc} \uparrow & \uparrow & \uparrow \\ \cos(\pi/4) & \cos(\pi/6) & \cos(2\pi/3) \end{array}$$

$$\rightarrow \cos(\pi/4) (\cos(\pi/6) + \cos(2\pi/3))$$

$$a = 1/6, \quad b = 2/3$$

$$(4) a) i) (z - 2i)^* = (x + yi - 2i)^* \\ = (x + yi + 2i)$$

$$ii) x - yi + 2i = 4iz + 3$$

$$\rightarrow x - yi + 2i = 4i(x + yi) + 3$$

$$\rightarrow x - yi + 2i = 4xi - 4y + 3$$

$$\boxed{\text{REAL}} \quad x = -4y + 3$$

$$\boxed{\text{IMAG}} \quad -y + 2 = 4x$$

$$\text{Real} \times 4 \rightarrow 4x = -16y + 12$$

$$\text{Equation: } -16y + 12 = -y + 2$$

$$\rightarrow 10 = 15y \rightarrow y = 10/15 = 2/3$$

$$x = -4(2/3) + 3 = 1/3$$


$$\Rightarrow 2 = \frac{1}{3} + \frac{2}{3}i$$

b) Sum of roots =  $-10i$

$$(p + qi) + (p - qi) = 2p$$

$\therefore p$  cannot be real if  $p - qi$  is a root

(5) a) i)



$$\textcircled{1} \quad y = 2(1+h)^2 - 5(1+h)$$

$$\Rightarrow y = 2(1 + 2h + h^2) - 5 - 5h$$

$$\Rightarrow y = 2 + 4h + 2h^2 - 5 - 5h$$

$$\Rightarrow y = 2h^2 - h - 3$$

$$\text{Gradient} = \frac{2h^2 - h - 3 - (-3)}{h}$$

$$= 2h - 1$$

ii) As  $h \rightarrow 0$ , gradient of tangent  $\rightarrow 2(0) - 1 = -1$

Gradient of  $x + y = 0 = -1$

$\therefore$  Lines are parallel

$$\begin{aligned} \text{b) } I &= \int_1^m 2x^{-2} - 5x^{-3} = \left[ -2x^{-1} + \frac{5}{2}x^{-2} \right]_1^m \\ &= \left[ \frac{-2}{x} + \frac{5}{2x^2} \right]_1^m = \left( \frac{-2}{m} + \frac{5}{2m^2} \right) - \left( \frac{-2}{1} + \frac{5}{2} \right) \\ &= \frac{-2}{m} + \frac{5}{2m^2} - 0.5 \end{aligned}$$

As  $m \rightarrow \infty$ ,  $\frac{-2}{m} \approx \frac{5}{2m^2} \rightarrow 0$

$\therefore I \rightarrow -0.5$

(6) a)  $\alpha + \beta = -3/2$        $\alpha\beta = -6/2 = -3$

b)  $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$

$$= \left(-\frac{3}{2}\right)^3 - 3(-3)\left(-\frac{3}{2}\right)$$

$$= -\frac{27}{8} - \frac{27}{2} = -\frac{135}{8}$$

$$\boxed{\text{Sum}} \quad \alpha + \frac{\alpha}{\beta^2} + \beta + \frac{\beta}{\alpha^2}$$

$$= \alpha + \beta + \cancel{\frac{\alpha^3}{\beta^2}} + \frac{\alpha^3}{\alpha^2 \beta^2} + \frac{\beta^2}{\alpha^2 \beta^2}$$

$$= (\alpha + \beta) + \frac{\alpha^3 + \beta^3}{(\alpha\beta)^2}$$

$$= -3/2 + \frac{-135/8}{(-3)^2} = -27/8$$

$$\boxed{\text{Product}} \quad \left( \alpha + \frac{\alpha}{\beta^2} \right) \left( \beta + \frac{\beta}{\alpha^2} \right) = \alpha\beta + \frac{\alpha\beta}{\alpha^2} + \frac{\alpha\beta}{\beta^2} + \frac{\alpha\beta}{\alpha^2 \beta^2}$$

$$= \alpha\beta + \frac{\beta}{\alpha} + \frac{\alpha}{\beta} + \frac{1}{\alpha\beta}$$

$$= \alpha\beta + \frac{\alpha^2 + \beta^2}{\alpha\beta} + \frac{1}{\alpha\beta}$$

$$= \alpha\beta + \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} + \frac{1}{\alpha\beta}$$

$$= -3 + \frac{(-3/2)^2 - 2(-3)}{-3} + \frac{1}{-3}$$

$$= -73/12$$

$$x^2 - \boxed{\text{Sum}} x + \boxed{\text{Product}} = 0$$

$$\Rightarrow x^2 + 27/8 x - 73/12 = 0$$

$$\times 24 \Rightarrow 24x^2 + 81x - 146 = 0$$

$$(7) \quad a) \quad \text{Let } f(x) = 4x^3 - 5x - 540,000$$

$$f(51) = 4(51)^3 - (51) - 540,000 = -9447$$

$$f(52) = 4(52)^3 - (52) - 540,000 = 22,380$$

Sign change,  $\therefore$  root lies between 51 & 52

$$b) \quad i) \quad S_n = \sum 4r^2 - 4r + 1$$

$$= 4 \sum r^2 - 4 \sum r + \sum 1$$

$$= \frac{1}{3} n (n+1) (2n+1) - 2n(n+1) + n$$

$$= \frac{1}{3} [ 2(n+1)(2n+1) - 6(n+1) + 3 ]$$

$$= \frac{1}{3} [ 2(2n^2 + n + 2n + 1) - 6n - 6 + 3 ]$$

$$= \frac{1}{3} [ 4n^2 + 6n + 2 - 6n - 6 + 3 ]$$

$$= \frac{1}{3} [ 4n^2 - 1 ] \quad \Rightarrow k = 4$$

$$\text{ii) } 6S_n = 2n [4n^2 - 1]$$

$$= 2n (2n+1) (2n-1)$$

$$= (2n-1) 2n (2n+1)$$

= product of 3 consecutive integers.

$$\text{c) let odd numbers be } 2n-1$$

$$\begin{array}{l} \text{eg } n = \{ 1 \quad 2 \quad 3 \quad 4 \quad \dots \\ \text{number} = \{ 1 \quad 3 \quad 5 \quad 7 \quad \dots \end{array}$$

$$\therefore \sum_{i=1}^n (2n-1)^2 > 180,000$$

$$\Rightarrow \frac{1}{3} (4n^2 - 1) > 180,000$$

$$\text{Try values: } n = 51 \rightarrow 176,851 \quad (\text{too small})$$

$$n = 52 \rightarrow 187,400 \quad (\text{too big})$$

$$\therefore n = 52 \text{ is smallest value of } n.$$

$$8) \text{ a) Stretch, SF 3 in } y\text{-direction}$$

$$\Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\text{b) i) } y = \sqrt{3}x \Rightarrow y = (\tan 60^\circ)x$$

$$\Rightarrow \begin{bmatrix} \cos(120^\circ) & \sin(120^\circ) \\ \sin(120^\circ) & -\cos(120^\circ) \end{bmatrix} = \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ \sqrt{3}/2 & 1/2 \end{bmatrix}$$

ii) New Matrix (2) x Matrix (1)

$$\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ \sqrt{3}/2 & 1/2 \end{bmatrix}$$

(9) a) Denominator =  $(x-3)(x+1)$

$\rightarrow x=3$  &  $x=-1$  are asymptotes

As  $x \rightarrow \infty$ ,  $y \rightarrow 1/1 \rightarrow y=1$  is asymptote

b) i)  $y = K \rightarrow K = \frac{x^2 - 2x + 1}{x^2 - 2x - 3}$

$\rightarrow K = \frac{(x-1)(x-1)}{(x-3)(x+1)}$  NOT NEEDED!  
 $\downarrow$

$\rightarrow K(x^2 - 2x - 3) = x^2 - 2x + 1$

$\rightarrow Kx^2 - 2Kx - 3K = x^2 - 2x + 1$

$\rightarrow Kx^2 - x^2 - 2Kx + 2x - 3K - 1 = 0$

$\rightarrow (K-1)x^2 - 2(K-1)x - (3K+1) = 0$

ii) Real roots  $\rightarrow b^2 - 4ac \geq 0$

$\rightarrow [-2(K-1)]^2 + 4(K-1)(3K+1) \geq 0$

$\rightarrow 4[K^2 - 2K + 1] + 4[3K^2 + K - 3K - 1] \geq 0$


(÷4)  $\rightarrow K^2 - 2K + 1 + 3K^2 - 2K - 1 \geq 0$

$\rightarrow 4K^2 - 4K \geq 0$

(÷4)  $\rightarrow K^2 - K \geq 0$

iii) At stationary points,  $K^2 - K = 0$

$\rightarrow K(K-1) = 0$



$\rightarrow K=1$

$K=0$

$\rightarrow y=1$  = Asymptote!

$\rightarrow y=0$

So  $y=1$  NOT stationary point

when  $y \neq 0$ ,  $x \rightarrow 0 \leq \frac{x^2 - 2x + 1}{x^2 + 3x - 3}$

$\rightarrow 0 = (x-1)(x-1)$

$\rightarrow x = 1$

$\therefore$  Only 1 stationary point at  $(1, 0)$

c)

